

other. *F. ulkei* is a boreal form whose main range is in the northern tier of states and southern Canada.

## LITERATURE CITED

CREIGHTON, W. S.

1950. The ants of North America. Bull. Mus. Comp. Zool., 104:1-585.

EMERY, C.

1893. Beiträge zur Kenntnis der nordamerikanischen Ameisen-fauna. Zool. Jahrb. Syst., 7:633-682.

WHEELER, W. M.

1913. A revision of the ants of the genus *Formica*. Bull. Mus. Comp. Zool., 53:379-565.

---

CORRECTION TO THE SYNONYMY OF THE ANT *CAMPONOTUS FORMOSENSIS* WHEELER.—In a joint paper (Yasumatsu and Brown, 1951, Jour. Fac. Agr., Univ. Kyushu, 10, cf. p. 42), the synonymy of *Camponotus formosensis* is cited incorrectly. Due to an unfortunate clerical error, the name *Camponotus maculatus taylori* var. *formosae* Wheeler and the variant nomenclatorial equivalents *C. barbatus* var. *formosae*, Emery and *C. barbatus taylori* var. *formosae*, Teranishi were wrongly included in this formal list of synonyms and must now be removed. The variety *formosae*, whatever may be its relationship to *maculatus* or *barbatus*, is of course not at all closely related to *C. formosensis* or any other member of the *C. herculeanus* group.—W. L. BROWN, JR., Museum of Comparative Zoology, Harvard University.